

Planar Embeddings with a Globally Attracting Fixed Point

Begoña Alarcón ^{*}, Víctor Guíñez [†] and Carlos Gutierrez [‡]

Abstract

We consider sufficient conditions which guarantee that an embedding from the plane $\mathbb{R}^2$ into itself has a unique fixed point. We study sufficient conditions which imply the appearing of a globally attracting fixed point for such an embedding.

1 Introduction

This work deals with embeddings from the plane $\mathbb{R}^2$ into itself. It was motivated by questions about the existence of unique fixed points, as well as questions about stability. (See [2], [4] [5], [11] and [13].) It was, in particular, inspired by the following:

DMY Question (Discrete Markus-Yamabe Question) [4, 5] Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a C^1 -map such that $f(0) = 0$ and $\text{Spec}(f) \subset B(0, 1)$. Is 0 a global attractor for the discrete dynamical system generated by f ?

To state our results, we shall need the following definitions.

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a differentiable map. Denote by $\text{Spec}(f)$ the set of eigenvalues of Df_p , for all $p \in \mathbb{R}^n$.

^{*}Work partially supported in part by MECD Grant # HBE2002-0006, and by MECD Grant # MTM2004-03244.

[†]Supported in part by FONDECYT Grant # 1050036.

[‡]The third author was supported in part by FAPESP Grant # TEMÁTICO 03/03107-9, by CNPq Grant # 306992/2003-5 and Project CAPES-MECD grant 071/04 and HBP2003-0017.

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a continuous map. We say that γ is an f -invariant ray if γ is a smooth embedded curve with no self-intersections starting at 0 and going to infinity such that $f(\gamma) \subseteq \gamma$. It is assumed that $0 \in \gamma$.

Let $\text{Emb}(\mathbb{R}^2)$ denote the set of topological embeddings $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$. We will denote by $\text{Emb}_+(\mathbb{R}^2)$ the subset of the orientation preserving maps of $\text{Emb}(\mathbb{R}^2)$ and by $\text{Emb}^d(\mathbb{R}^2)$ the subset of differentiable maps of $\text{Emb}(\mathbb{R}^2)$. Finally, let $\text{Emb}_+^d(\mathbb{R}^2) = \text{Emb}^d(\mathbb{R}^2) \cap \text{Emb}_+(\mathbb{R}^2)$.

Let $f \in \text{Emb}(\mathbb{R}^2)$ and let $p \in \mathbb{R}^2$. We define the ω -limit set of p , $\omega(p)$, the set (possibly empty) of points $x \in \mathbb{R}^2$ for which there exists a sequence of positive integers $n_k \rightarrow \infty$ such that $f^{n_k}(p) \rightarrow x$ as $k \rightarrow \infty$.

We will say that $x \in \mathbb{R}^2$ is a non-wandering point of $f \in \text{Emb}(\mathbb{R}^2)$ if for every neighborhood U of x , there exist an integer $n > 2$ and a point $q \in U$ such that $f^n(q) \in U$. We denote by $\Omega(f)$ the set of non-wandering points of f .

Let $f \in \text{Emb}(\mathbb{R}^2)$

- We say that $0 \in \mathbb{R}^2$ is a local attractor (resp. a local repeller) for f if there exists a compact disc D , contained in the domain of definition of f (resp. contained in the domain of definition of f^{-1}), which is a neighborhood of 0 such that $f(D) \subset \text{Int}(D)$ (resp. $f^{-1}(D) \subset \text{int}(D)$) and $\bigcap_{n=1}^{\infty} f^n(D) = \{0\}$ (resp. $\bigcap_{n=1}^{\infty} f^{-n}(D) = \{0\}$).

Note that if $f \in \text{Emb}^1(\mathbb{R}^2)$ has a hyperbolic attractor at 0, our definition coincides with the classical one [8].

- We shall say that $0 \in \mathbb{R}^2$ is a global attractor for f if 0 is a local attractor and for all $p \in \mathbb{R}^2$, $\omega(p) = \{0\}$.

If $f : \mathbb{R}^2 \cup \{\infty\} \rightarrow \mathbb{R}^2 \cup \{\infty\}$ is a homeomorphism of the Riemann Sphere, with $f(\infty) = \infty$, we may similarly define when ∞ is either an attractor or a repeller.

A compact set $W \subset \mathbb{R}^2$ is a window of $f \in \text{Emb}(\mathbb{R}^2)$ if for all $p \in \mathbb{R}^2$ there exists a $n_0 \in \mathbb{N}$ such that $f^{n_0}(p) \in W$. We will say that $f \in \text{Emb}(\mathbb{R}^2)$ is a dissipative map if there exists a window for f .

Assume that $f \in \text{Emb}(\mathbb{R}^2)$ such that $f(p) = p$. The fixed point p is stable in the sense of Lyapunov (or Lyapunov stable) if every neighborhood U of

p contains another neighborhood V such that, for each $n \geq 0$, $f^n(V)$ is well defined and $f^n(V) \subset U$.

It was proved in [4] that the DMY Question had a positive answer for polynomial diffeomorphisms of $\mathbb{R}^2$ but was false even for rational diffeomorphism of $\mathbb{R}^2$. Therefore, we wondered if the DMY Question had a positive answer, for smooth diffeomorphisms of $\mathbb{R}^2$ under the additional assumption

(a) ∞ is a repeller.

We show that this assumption is not good enough either: there exists a smooth diffeomorphism $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ having an order four periodic point and such that ∞ is a repeller, $f(0) = 0$ and $\text{Spec}(f) \subset B(0, 1)$. (See Theorem 4.4).

Since every global diffeomorphism f of $\mathbb{R}^2$ having 0 as a global attractor satisfies

(b) f has an f -invariant ray

(see Proposition 5.2 where we show that for such an f , there is a foliation of $\mathbb{R}^2 \setminus 0$ by f -invariant rays), we study the MYD Question under the additional assumptions (a) and/or (b) obtaining the following five results:

Theorem 2.8. Let $f \in \text{Emb}_+^d(\mathbb{R}^2)$ be such that $f(0) = 0$, and that it has an f -invariant ray γ . If $\text{Spec}(f) \cap [1, 1 + \epsilon) = \emptyset$, for some $\epsilon > 0$, then $\Omega(f) \subset \gamma$ and either $\omega(p) = \{0\}$ or $\omega(p) = \emptyset$, for all $p \in \mathbb{R}^2$.

Theorem 2.9 Let $f \in \text{Emb}_+^d(\mathbb{R}^2)$ be such that $f(0) = 0$, and that it has an f -invariant ray γ . If $\text{Spec}(f) \subset B(0, 1)$, then 0 is a local attractor and $\Omega(f) = \{0\}$.

Theorem 2.11 Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a differentiable map such that $f(0) = 0$. Then:

- (a) If $\text{Spec}(f) \cap ([1, 1 + \epsilon) \cup [0, \epsilon)) = \emptyset$, for some $\epsilon > 0$, then f is an embedding with $\text{Fix}(f) = \{0\}$.
- (b) If $\text{Spec}(f) \subset B(0, 1) \setminus [0, \epsilon)$, for some $\epsilon > 0$, then not only is f an embedding with $\text{Fix}(f) = \{0\}$, and 0 is a local attractor for f .

Relevant to our results, Figure 1 shows the phase portrait of a smooth flow transversal to the unit circle ∂D such that, the time-one-map $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$,

induced by the flow, which can be assumed to be a smooth diffeomorphism of $\mathbb{R}^2$, satisfies $f(D) \subset \text{Int}(D)$ (where D is the unit disc.) and, for all $p \in \mathbb{R}^2$, $\omega(p) = \{0\}$. Observe that any trajectory of the flow meeting ∂D is an f -invariant ray, while the origin 0 is not a local attractor. For further examples, see [2].




Figure 1: A vector field which does not have the origin as an attractor.

Theorem 3.3. Let $f \in \text{Emb}(\mathbb{R}^2)$ be a dissipative map such that $f(0) = 0$, and that it has an f -invariant ray γ . Under either of the following conditions

- a) $f \in \text{Emb}_+(\mathbb{R}^2)$ and $\text{Fix}(f) = \{0\}$,
- b) $\text{Fix}(f^2) = \{0\}$,

we have that $\omega(p) = \{0\}$, for all $p \in \mathbb{R}^2$. Moreover, if 0 is a locally Lyapunov stable fixed point, then 0 is a global attractor for f .

Theorem 3.6. Let $f \in \text{Emb}(\mathbb{R}^2)$ be of class C^1 such that $f(0) = 0$, and that it satisfies the following three conditions:

- (i) There exist real numbers $R > 0$ and $0 < \alpha < 1$ such that $\|Df_p \cdot p\| < \alpha \|p\|$, for every $\|p\| > R$.
- (ii) $\text{Spec}(f) \subset B(0, 1)$.
- (iii) There exists a f -invariant ray γ .

Under either of the following two conditions:

- (1) f is orientation preserving

$$(2) \text{ Spec}(f) \cap \mathbb{R} = \emptyset,$$

we have that 0 is a global attractor for f .

In obtaining our results, we have profited from the main results of [1], [3], [6], [10], [14].

Section 2 deals with conditions which imply that a continuous map from $\mathbb{R}^2$ into itself is an embedding and/or has a unique fixed point. In Section 3, we study dissipative embeddings having a global attractor. Some important examples are considered in Section 4. Section 5 presents the results which motivated our assumptions.

2 Embeddings with only one fixed point

We shall need the following

Theorem 2.1 (Murthy [10], 1998). *Let $f \in \text{Emb}_+(\mathbb{R}^2)$. If f has a periodic point p , then the set $\text{Fix}(f)$ of the fixed points of f is nonempty.*

Corollary 2.2. *Let $f \in \text{Emb}_+(\mathbb{R}^2)$. If $\Omega(f) \neq \emptyset$ then $\text{Fix}(f) \neq \emptyset$.*

Proof. Suppose by contradiction that $\text{Fix}(f) = \emptyset$. By Theorem 2.1, $\Omega(f)$ does not contain periodic points. Let $p \in \Omega(f)$. Then there exist a disc U , an integer $n > 2$ and a point $q \in U$ such that $f^n(q) \in U$, and for all, $k = 1, 2, \dots, n-1$, $U \cap f^k(U) = \emptyset$. Let $g : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be an orientation-preserving embedding which is a perturbation of f supported in U , such that $g(f^n(q)) = f(q)$. Then q is periodic point of g of period n . Therefore, by Theorem 2.1, g has a fixed point q_0 . Certainly $q_0 \notin U$ and so $q_0 \in \text{Fix}(f)$. This contradiction proves the result. $\square$

Corollary 2.3. *If $U \subset \mathbb{R}^2$ is an open set homeomorphic to $\mathbb{R}^2$ and $g : U \rightarrow U$ is an orientation preserving embedding with $\Omega(g) \neq \emptyset$, then g has a fixed point.*

Proof. If $H : U \rightarrow \mathbb{R}^2$ is a homeomorphism then we may apply Corollary 2.2 to $f = H \circ g \circ H^{-1}$. $\square$

We remark that the version of Theorem 2.1 for chain-recurrent-points can be found in [10].

Theorem 2.4 (Fernandes, Gutierrez and Rabanal [6], 2004). *Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a differentiable (not necessarily C^1) map. If for some $\epsilon > 0$, $\text{Spec}(f) \cap [0, \epsilon) = \emptyset$, then f is an embedding.*

Corollary 2.5. *Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a differentiable map such that $\text{Spec}(f) \cap [1, 1 + \epsilon) = \emptyset$, for some $\epsilon > 0$. Then $\text{Fix}(f)$ is either empty or a one-point-set.*

Proof. Let $g = f - \text{Id}$, where $\text{Id} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is the identity map. Certainly $\lambda \in \text{Spec}(f)$ if and only if $\lambda - 1 \in \text{Spec}(g)$. So there exists an $\epsilon > 0$ such that $\text{Spec}(g) \cap [0, \epsilon) = \emptyset$. By Theorem 2.4, g is injective; this implies the conclusion. $\square$

The following result gives us a condition for a differentiable map be an embedding having a unique fixed point:

Corollary 2.6. *Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a differentiable map such that $f(0) = 0$ and $\text{Spec}(f) \cap ([0, \epsilon) \cup [1, 1 + \epsilon)) = \emptyset$, for some $\epsilon > 0$. Then $f \in \text{Emb}^d(\mathbb{R}^2)$ such that $\text{Fix}(f) = \{0\}$.*

Proof. It follows immediately from Theorem 2.4 and Corollary 2.5. $\square$

Lemma 2.7. *Let $f \in \text{Emb}(\mathbb{R}^2)$ with $f(0) = 0$ and having an f -invariant ray γ . Under either of the following conditions*

- a) $f \in \text{Emb}_+(\mathbb{R}^2)$ and $\text{Fix}(f) = \{0\}$,
- b) $\text{Fix}(f^2) = \{0\}$,

we have that $\Omega(f) \subset \gamma$ and either $\omega(p) = \{0\}$ or $\omega(p) = \emptyset$, for all $p \in \mathbb{R}^2$.

Proof. Suppose a) is satisfied. If 0 is the only fixed point of f and there exists a f -invariant ray γ , then $f : \mathbb{R}^2 \setminus \gamma \rightarrow \mathbb{R}^2 \setminus \gamma$ is a fixed-point-free orientation preserving embedding.

As $\mathbb{R}^2 \setminus \gamma \subset \mathbb{R}^2$ is an open set homeomorphic to $\mathbb{R}^2$, by Corollary 2.2, $\Omega(f|_{\mathbb{R}^2 \setminus \gamma}) = \emptyset$. Then $\Omega(f) \subset \gamma$ and since $f|_\gamma : \gamma \rightarrow \gamma$ is a one dimensional homeomorphism with only one fixed point, then given $p \in \mathbb{R}^2$ either $\omega(p) = \{0\}$ or $\omega(p) = \emptyset$.

If b) is satisfied we may apply a) to f^2 , which is always orientation preserving, and obtain the required conclusion. $\square$

Theorem 2.8. *Let $f \in \text{Emb}_+^d(\mathbb{R}^2)$ be such that $f(0) = 0$, and that it has an f -invariant ray γ . If $\text{Spec}(f) \cap [1, 1 + \epsilon) = \emptyset$, for some $\epsilon > 0$, then $\Omega(f) \subset \gamma$ and either $\omega(p) = \{0\}$ or $\omega(p) = \emptyset$, for all $p \in \mathbb{R}^2$.*

Proof. It follows directly from Lemma 2.7 and Corollary 2.5. $\square$

Theorem 2.9. *Let $f \in \text{Emb}_+^d(\mathbb{R}^2)$ be such that $f(0) = 0$, and that it has an f -invariant ray γ . If $\text{Spec}(f) \subset B(0, 1)$, then 0 is a local attractor and $\Omega(f) = \{0\}$.*

Proof. It follows from the Hartman and Grobman Theorem [8] that 0 is a local attractor. By applying Theorem 2.8, we obtain this result. $\square$

Remark 2.10. *Under the condition of Theorem 3.3, if 0 is stable in the sense of Lyapunov, we have $\text{index}(f, 0) = 1$ ([11]).*

Theorem 2.11. *Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a differentiable map such that $f(0) = 0$. Then:*

- (a) *If $\text{Spec}(f) \cap ([1, 1 + \epsilon) \cup [0, \epsilon)) = \emptyset$, for some $\epsilon > 0$, then f is an embedding with $\text{Fix}(f) = \{0\}$.*
- (b) *If $\text{Spec}(f) \subset B(0, 1) \setminus [0, \epsilon)$, for some $\epsilon > 0$, then not only is f an embedding with $\text{Fix}(f) = \{0\}$, and 0 is a local attractor for f .*

Proof. Item (a) follows from Corollary 2.5. Item (b) follows from (a) and from Hartman and Grobman Theorem ([8]). $\square$

The proof of theorem below can be found in [1].

Theorem 2.12. *Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a differentiable map with $f(0) = 0$. If $\text{Spec}(f) \cap \mathbb{R} = \emptyset$, then f is an embedding with $\text{Fix}(f^2) = \{0\}$.*

3 Dissipative embeddings and attractors

We shall need the following result (see also [9]):

Proposition 3.1. *(Richeson and Wiseman [12], 2002). Let X be a metric locally compact set and $f : X \rightarrow f(X)$ be an homeomorphism. The following are equivalent:*

- (i) f is a dissipative map.
- (ii) For all compact set $D \subset X$ there exists a window W for f such that $D \subset \text{int}(W)$, $f(W) \subset W$ and $\forall p \in X \exists n_0$ such that $f^{n_0}(p) \in \text{int}(W)$.
- (iii) For all compact set $D \subset X$ there exists a window W for f such that $D \subset \text{int}(W)$ and $f(W) \subset \text{int}(W)$.

Remark 3.2. *It follows from Proposition right above that if $f \in \text{Emb}(\mathbb{R}^2)$ is a dissipative embedding, then there exists a compact set D such that for all $p \in \mathbb{R}^2$ the set $\omega(p)$ is non empty and $\omega(p) \subset D$. Note that if $f(0) = 0$, then $0 \in D$.*

Theorem 3.3. *Let $f \in \text{Emb}(\mathbb{R}^2)$ be a dissipative map such that $f(0) = 0$, and that it has an f -invariant ray γ . Under either of the following conditions*

- a) $f \in \text{Emb}_+(\mathbb{R}^2)$ and $\text{Fix}(f) = \{0\}$,
- b) $\text{Fix}(f^2) = \{0\}$,

we have that $\omega(p) = \{0\}$, for all $p \in \mathbb{R}^2$. Moreover, if 0 is a locally Lyapunov stable fixed point, then 0 is a global attractor for f .

Proof. As f is a dissipative map, there exists a compact set $D \subset \mathbb{R}^2$ such that for all $p \in \mathbb{R}^2$, $\omega(p) \neq \emptyset$ and $\omega(p) \subset D$. Using Lemma 2.7 we conclude that for all $p \in \mathbb{R}^2$, $\omega(p) = \{0\}$. Therefore, if 0 is locally stable in the sense of Lyapunov, we may obtain that 0 is a global attractor. $\square$

Remark 3.4. *Under the condition of Theorem 3.3, if 0 is stable in the sense of Lyapunov, we have $\text{index}(f, 0) = 1$ ([11]).*

Lemma 3.5. *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a C^1 -map such that $f(0) = 0$. Suppose that there exist $R > 0$ and $0 < \alpha < 1$ such that*

$$\forall \|p\| > R, \quad \|Df_p \cdot p\| < \alpha \|p\|.$$

There exist $S_0 > R$ and $\alpha < \mu < 1$ such that if $p \in \mathbb{R}^n$ and $\|p\| \geq S_0$ then $\|f(p)\| \leq \mu \|p\|$.

Proof. As $\bar{B}(0, R)$ is a compact set, there exists a real number $M > 1$ such that $\|Df_q\| \leq M$ for all $q \in \bar{B}(0, R)$.

Let $\mu = \frac{\alpha+1}{2}$ and $S_0 = \frac{2(MR-\alpha R)}{1-\alpha}$. Notice that $0 < \alpha < \mu < 1$ and $S_0 > R$. Given $p \in \mathbb{R}^n$ with $\|p\| \geq S_0$, let $\gamma : [0, \|p\|] \rightarrow \mathbb{R}^n$ be defined by $\gamma(t) = (tp)/\|p\|$. Then

$$\begin{aligned} \|f(p)\| &\leq \int_0^R \|(f \circ \gamma)'(t)\| dt + \int_R^{\|p\|} \|(f \circ \gamma)'(t)\| dt \\ &\leq MR + \alpha (\|p\| - R) \\ &= \frac{1-\alpha}{2} \cdot \frac{2(MR-\alpha R)}{1-\alpha} + \alpha \|p\| \\ &\leq \frac{1-\alpha}{2} \|p\| + \alpha \|p\| = \mu \|p\|. \end{aligned}$$

□

Theorem 3.6. *Let $f \in \text{Emb}(\mathbb{R}^2)$ be of class C^1 such that $f(0) = 0$, and that it satisfies the following three conditions:*

- (i) *There exist real numbers $R > 0$ and $0 < \alpha < 1$ such that $\|Df_p \cdot p\| < \alpha \|p\|$, for every $\|p\| > R$.*
- (ii) *$\text{Spec}(f) \subset B(0, 1)$.*
- (iii) *There exists a f -invariant ray γ .*

Under either of the following two conditions:

- (1) *f is orientation preserving*
- (2) *$\text{Spec}(f) \cap \mathbb{R} = \emptyset$,*

we have that 0 is a global attractor for f .

Proof. Suppose that (1) is satisfied. By item (i) and Lemma 3.5, f is dissipative. Therefore, by items (ii), (iii) and Hartman and Grobman Theorem [8], we may apply Theorem 2.9 to obtain the requested conclusion: 0 is a global attractor for f .

By Theorem 2.12, the proof of this theorem under assumption (2) is the same as that under the assumption (1); notice that f^2 is always orientation preserving. $\square$

4 Examples

The following two examples can be found in [4].

Theorem 4.1 (Szlenk's Example). *Let $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be defined by*

$$F(x, y) = \left(-\frac{ky^3}{1+x^2+y^2}, \frac{kx^3}{1+x^2+y^2} \right), \quad \text{where } k \in \left(1, \frac{2}{\sqrt{3}}\right).$$

The map F satisfies the following properties:

1. *Set $p = (x, y) \in \mathbb{R}^2$ and let λ be an eigenvalue of $JF(p)$. If $xy = 0$ then $\lambda = 0$. Otherwise $\lambda \notin \mathbb{R}$ and $|\lambda| < \sqrt{3}k/2$.*
2. *$F^4((k-1)^{-1/2}, 0) = ((k-1)^{-1/2}, 0)$.*
3. *F is injective.*

Theorem 4.2 (Szlenk–Cima–Gasull–Mañosas's Example). *Let $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be as in theorem 4.1 and let $0 < a < 1$. Let $G_a : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be defined by*

$$G_a(x, y) = F(x, y) - a(x, y).$$

If a is small enough, the map G_a is a global diffeomorphism of $\mathbb{R}^2$ onto itself which satisfies the following properties:

- (a) *for all $x \in \mathbb{R}^2$, $\text{Spec}(G_a)(x) \subset B(0, 1)$*
- (b) *$G_a(0) = 0$ and there exists $p \in \mathbb{R}^2 \setminus \{0\}$ such that $(G_a)^4(p) = p$.*

Remark 4.3. For the Szlenk's map F , there are points p such that $\lim_{n \rightarrow \infty} F^n(p) = \infty$ and therefore ∞ is not a repellor. Also there is numerical evidence that for the Szlenk–Cima–Gasull–Mañosas's map G_a , ∞ is not a repellor.

The next example shows that DMY Question has a negative answer, for smooth diffeomorphisms of $\mathbb{R}^2$, under the additional condition that ∞ is a repellor.

Theorem 4.4. *There exists a smooth diffeomorphism $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ having an order four periodic point and such that ∞ is a repellor, $f(0) = 0$ and $\text{Spec}(f) \subset B(0, 1)$.*

We introduce some notations and Lemmas that we will use in the proof of Theorem 4.4.

Notation 4.5. If $A : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear map, we denote by $\text{sr}(A)$ the spectral radius of A :

$$\text{sr}(A) = \max \{ |\lambda| : \lambda \text{ is an eigenvalue of } A \}$$

As usual, the norm $\|A\|$ of A is

$$\|A\| = \sup \{ \|Av\| : \|v\| = 1 \}.$$

It is well known that

$$\text{sr}(A) \leq \|A\|.$$

If $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a diffeomorphism we denote by

$$\text{sr}(F) = \sup \{ \text{sr}(DF_{(p)}) : p \in \mathbb{R}^2 \}.$$

Lemma 4.6. *Given $R > 0$, $C > 0$ and $0 < \varepsilon < \frac{1}{8C}$, there exists a smooth function*

$$\phi : [0, \infty) \rightarrow \left[\frac{1}{2C}, 1 \right]$$

which verifies

- 1) $\phi(r) = 1$, for all $r \in [0, R]$,
- 2) $\phi'(r) \leq 0$ and $|\phi'(r) \cdot r| < \varepsilon$ for all $r \in [0, \infty)$,

3) there is an integer N such that $\phi(r) = \frac{1}{2C}$, for all $r > N$.

Proof. First we define $\phi(r) = 1$, for all $r \in [0, R]$. Hence, we consider the sequence $(S(n))$ defined for all positive integer n by

$$S(n) = 1 - \frac{\varepsilon}{8} \left(\frac{1}{R+1} + \frac{1}{R+2} + \cdots + \frac{1}{R+n} \right).$$

Observe that $S(n) \rightarrow -\infty$ as $n \rightarrow \infty$. Let N be the integer such that $S(n) > \frac{1}{2C}$ for $n = 1, \dots, N-1$, and $S(N) \leq \frac{1}{2C}$.

For all positive integer $n \geq 1$ we define

$$\begin{cases} \phi(R+n) = S(n), & \text{if } 1 \leq n \leq N-1, \\ \phi(R+n) = \frac{1}{2C}, & \text{if } n \geq N. \end{cases}$$

For each integer n such that $1 \leq n \leq N-1$ we consider a smooth map $\phi : [R+n, R+n+1] \rightarrow [S(n+1), S(n)]$ which is flat at $R+n$ and $R+n+1$ and defined as above and such that

$$0 \leq -\phi'(r) = |\phi'(r)| \leq \frac{2(S(n) - S(n+1))}{(R+n+1) - (R+n)} = \frac{\varepsilon}{4(R+n+1)}.$$

Finally, for $r > R+N$ we define $\phi(r) = \frac{1}{2C}$.

Then the map ϕ verifies the conditions because, for $n = 1, \dots, N$ and $R+n-1 \leq r \leq R+n$ we have

$$|\phi'(r) \cdot r| \leq \frac{\varepsilon}{4(R+n)} \cdot (R+n) = \frac{\varepsilon}{4} < \varepsilon.$$

□

Lemma 4.7. Let $M_2(\mathbb{R})$ be the space of 2×2 real matrices and let C be a positive constant. Denote by $\mathcal{A}$ the compact set consisting of $A \in M_2(\mathbb{R})$ such that $\|A\| \leq C$ and $\text{sr}(A) \leq 0.9$. There exists $\frac{1}{8C} > \varepsilon > 0$ such that if

$$B = \begin{pmatrix} b + \varepsilon_1 & \varepsilon_2 \\ \varepsilon_3 & b + \varepsilon_4 \end{pmatrix}$$

satisfies: $\frac{1}{2C} \leq b \leq 1$, and $\max \{ |\varepsilon_1|, |\varepsilon_2|, |\varepsilon_3|, |\varepsilon_4| \} < \varepsilon$, then

$$\text{sr}(BA) \leq 0.95,$$

for all $A \in \mathcal{A}$.

Proof. Writting $B = bI + E$, we have $BA = bA + EA$ and

$$\det(BA - \lambda I) = b^2 \det \left(A + \frac{1}{b} EA - \frac{\lambda}{b} I \right),$$

which implies

$$\text{sr}(BA) = a \text{sr} \left(A + \frac{1}{b} EA \right).$$

Then, given $\delta > 0$ there exist $0 < \varepsilon < \frac{1}{8C}$ such that

$$\text{sr} \left(A + \frac{1}{b} EA \right) \leq \text{sr}(A) + \frac{\delta}{b},$$

that is, such that

$$\text{sr}(BA) \leq b \text{sr}(A) + \delta.$$

Hence, it sufficient to consider $0 < \delta < 0.95 - b \cdot 0.9$. $\square$

Proof of Theorem 4.4. Consider the Szlenk–Cima–Gasull–Mañosas’s Example

$$G_a(x, y) = \left(-\frac{ky^3}{1+x^2+y^2}, \frac{kx^3}{1+x^2+y^2} \right) - a(x, y),$$

with $k \in (1, \frac{2}{\sqrt{3}})$ and $0 < a < a_0 < 1$ such that G_a verifies Theorem 4.2. Also consider a constant $C > 0$ such that for all $(x, y) \in \mathbb{R}^2$,

$$\|D G_a(x, y)\| \leq C. \quad (1)$$

Since, $\text{sr}(G_0) < \sqrt{3}k/2$ and $\text{Spec}(G_a)(x, y) = \text{Spec}(G_0)(x, y) - a$, for all $(x, y) \in \mathbb{R}^2$, there exist $0 < a_0 < 1$ and $1 < k_0 < \frac{2}{\sqrt{3}} \cdot (0, 88) (\approx 1, 01614)$, such that

$$\text{sr}(G_a) \leq 0.9, \quad \text{for all } 0 < a < a_0 \quad \text{and} \quad 1 < k < k_0. \quad (2)$$

Associated to the numbers $R = \frac{2}{\sqrt{k}-1}$, $C > 0$ as in (1) and $0 < \varepsilon < \frac{1}{8C}$ as in Lemma 4.7, consider the smooth function

$$\phi : [0, \infty) \rightarrow \left[\frac{1}{2C}, 1 \right]$$

of Lemma 4.6. We will prove that

$$f = H \circ G_a : \mathbb{R}^2 \rightarrow \mathbb{R}^2,$$

with a (and k) as in (2), and with H defined by

$$H(x, y) = (\phi(\sqrt{x^2 + y^2})x, \phi(\sqrt{x^2 + y^2})y),$$

verifies our Theorem.

1) To prove f is a smooth diffeomorphism we show that H is a smooth diffeomorphism. If $r = \sqrt{x^2 + y^2}$, we have

$$JH(x, y) = \begin{pmatrix} \phi(r) + \phi'(r) \cdot \frac{x^2}{r} & \phi'(r) \cdot \frac{xy}{r} \\ \phi'(r) \cdot \frac{xy}{r} & \phi(r) + \phi'(r) \cdot \frac{y^2}{r} \end{pmatrix}$$

Since

$$\phi(r) > \frac{1}{2C} > 4\varepsilon > \varepsilon > |\phi'(r) \cdot r|,$$

implies

$$\det DH(x, y) = \phi(r) (\phi(r) + \phi'(r) \cdot r) = \phi(r) (\phi(r) - |\phi'(r) \cdot r|) > 0,$$

and

$$\lim_{|(x, y)| \rightarrow \infty} |H(x, y)| = \infty,$$

the map H is a global diffeomorphism.

2) To prove $\text{sr}(f) < 1$ first observe that $DG_a(x, y)$ is in the set $\mathcal{A}$ of Lemma 4.6, for all (x, y) . Also, if $r = |Ga(x, y)|$ and $(u, v) = Ga(x, y)$ we have

$$\frac{1}{2C} \leq \phi(r) \leq 1,$$

and

$$\max\{|\phi'(r) \cdot \frac{u^2}{r}|, |\phi'(r) \cdot \frac{uv}{r}|, |\phi'(r) \cdot \frac{v^2}{r}|\} \leq |\phi'(r) \cdot r| < \varepsilon.$$

Then, by Lemma 4.7, we obtain

$$\text{sr}(f) \leq 0.95.$$

3) Clearly $f(0, 0) = (0, 0)$. Moreover, since $p = (R/2, 0)$ is a hyperbolic order four periodic point of G_0 , then not only the diffeomorphism G_a has a hyperbolic order four periodic point in the puncture ball $B(0, R) - \{0\}$ but also f .

4) Finally

$$|f(x, y)| = \frac{|G_a(x, y)|}{2C} \leq \frac{1}{2}|(x, y)|,$$

for all $|(x, y)| \geq R + N$ with N as in item 3) of Lemma 4.6. Therefore, ∞ is a repeller for f and the proof is complete. $\square$

5 Final remarks

The following result might have motivated the DMY Question:

Proposition 5.1. *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a C^1 -map such that $f(0) = 0$. If for all $p \in \mathbb{R}^n$, $\|Df_p\| < 1$, then 0 is a global attractor for f .*

Proof. Let $q \in \mathbb{R}^n$ and $S = \|q\|$. As f is a C^1 -map, and $\bar{B}(0, S)$ is a compact set there exists a real number $M > 0$ such that for all $p \in \bar{B}(0, S)$, $\|Df_p\| \leq M < 1$. Hence $f(\bar{B}(0, S)) \subseteq B(0, S)$ and so $f|_{\bar{B}(0, S)}$ is a M -contraction. Therefore $f^k(q) \rightarrow 0$ as $k \rightarrow \infty$. As $q \in \mathbb{R}^n$ is an arbitrary point, 0 is a global attractor for the discrete dynamical system generated by f . $\square$

Next proposition justifies one of our main assumptions.

Proposition 5.2. *Let $f : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0)$ be a C^r -diffeomorphism, with $r \geq 1$, having 0 as a global attractor. Then there exists a C^r -foliation of $\mathbb{R}^2 - \{0\}$ by f -invariant rays.*

Proof. Given $0 < r_1 < r_2$ and $r > 0$ let denote $A(r_1, r_2) = \{p \in \mathbb{R}^2 : r_1 \leq \|p\| \leq r_2\}$, $B(r) = \{p \in \mathbb{R}^2 : \|p\| < r\}$ and $S(r) = \{p \in \mathbb{R}^2 : \|p\| = r\}$. Also, given a simple closed curve $\gamma \subset \mathbb{R}^2$, we denote by γ^- (resp. γ^+) the union of γ with the bounded (resp. unbounded) connected component of $\mathbb{R}^2 - \gamma$.

Let D be a smooth compact disk such that $0 \in \text{int}(D)$, $f(D) \subset \text{int}(D)$ and $\bigcup_{n \in \mathbb{Z}} f^n(D) = \mathbb{R}^2$. Denote by $C = \partial D$ the boundary of D . Let $V(C)$ be a small smooth tubular neighborhood of C such that $f(V(C)) \cap V(C) = \emptyset$. For $\varepsilon > 0$ small, consider a smooth embedding

$$g_1 : A(2 - 2\varepsilon, 2 + 2\varepsilon) \rightarrow V(C) \quad \text{with} \quad g_1(S(2)) = C.$$

and define the smooth embedding $g_2 = f \circ g_1 \circ T : A(1 - \varepsilon, 1 + \varepsilon) \rightarrow (V(C))$, where $T(x, y) = 2(x, y)$. In this way,

- (1) restricted to $A(2 - 2\varepsilon, 2 + 2\varepsilon)$, the map $f \circ g_1$ equals to $g_2 \circ T^{-1}$.

We claim that

- (2) there exists a smooth embedding $H_0 : A(1 - \varepsilon, 2 + 2\varepsilon) \rightarrow \mathbb{R}^2$ which is an extension of both g_1 and g_2 .

In fact, let $G_1 : S(2 - 2\varepsilon)^+ \rightarrow g_1(S(2 - 2\varepsilon))^+$ and $G_2 : S(1 + \varepsilon)^- \rightarrow g_2(S(1 + \varepsilon))^-$ be smooth diffeomorphisms which are extensions of g_1 and g_2 , respectively (see Theorems 8.3.3 and 8.1.9 of [7]). Using Theorem 8.3.2 of [7] we can find a smooth diffeomorphism $H : \mathbb{R}_\infty^2 \rightarrow \mathbb{R}_\infty^2$ diffeotopic to the identity of $\mathbb{R}^2$ which restricted to $S(2 - 2\varepsilon)^+$ and $S(1 + \varepsilon)^-$ coincides with G_1 and G_2 , respectively. The proof of claim (2) is obtained by defining H_0 to be equal to the restriction of H to $A(1 - \varepsilon, 2 + 2\varepsilon)$.

On the annulus $A(1 - \varepsilon, 2 + \varepsilon)$ we consider the foliation $\mathcal{L}_0$ whose leaves are the linear segments $r_p = \{tp : 1 - \varepsilon \leq t \leq 2 + \varepsilon\}$ with $p \in S(1)$. We claim that

- (3) If $\ell \subset A(2 - 2\varepsilon, 2 + 2\varepsilon)$, is an arc of a leaf of $\mathcal{L}_0$, then both $H(\ell)$ and $f(H(\ell))$ are arcs of the same leaf of the foliation $H(\mathcal{L}_0)$.

In fact, using (1), we obtain that

$$f \circ H(\ell) = f \circ g_1(\ell) = g_2 \circ T^{-1}(\ell) = H(T^{-1}(\ell)).$$

The claim follows from this and from the fact that ℓ and $T^{-1}(\ell)$ are arcs of the same leaf of $\mathcal{L}_0$.

Then, by (2) and (3), $\mathcal{F} = \bigcup_{k \in \mathbb{Z}} f^k(H(\mathcal{L}_0))$ is a smooth foliation on $\mathbb{R}^2 - \{0\}$ by f -invariant rays.

□

References

- [1] B. Alarcón, C. Gutierrez and J. Martínez-Alfaro. *Planar injective maps and periodic orbits*. Em preparation.
- [2] N. P. Bhatia and G. P. Szeg'ó. *Stability Theory of Dynamical Systems*. Springer-Verlag, 2002.
- [3] P. L. Calvez and J.-C. Yoccoz. Un théorème d'indice pour les homéomorphismes du plan au voisinage d'un point fixe. *Annals of Mathematics*, 146:241–293, 1997.
- [4] A. Cima, A. Gasull, and F. Mañosas. The discrete markus-yamabe problem. *Nonlinear Analysis*, 35:343354, 1999.
- [5] A. van den Essen. Polynomial automorphisms and the jacobian conjecture. 190, 2000.
- [6] A. Fernandez, C. Gutierrez, and R. Rabanal. Global asymptotic stability for differentiable vector fields of $\mathbb{R}^2$. *J. Differential Equations*, 2004.
- [7] Morris W. Hirsch. *Differential Topology*. Graduate Texts in Mathematics 33, Springer–Verlag, 1976.
- [8] J. Palis and W. de Melo. *Geometric Theory of Dynamical Systems*. Springer Verlag, New York (1982).
- [9] R. F. Martins. *Dinámica de sistemas dissipativos e não-autónomos com uma coordenada angular*. PhD thesis, Universidad de Granada, 2005.
- [10] P. Murthy. Periodic solutions of two-dimensional forced systems: The massera theorem and its extension. *Journal of Dynamics and Differential Equations*, 10(2):275–302, 1998.
- [11] R.Ortega. Some applications of the topological degree to stability theory. *Topological Methods in Differential Equations and Inclusions*. (Montreal, PQ, 1994), 377–409, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 472, Kluwer Acad. Publ., Dordrecht, 1995.
- [12] D. Richeson and J. Wiseman. A fixed point theorem for bounded dynamical systems. *Illinois J. Math.*, 46:491–495, 2002.

- [13] R. C. Robinson. *An introduction to Dynamical Systems Continuous and Discrete*. Pearson Education, Inc, 2004.
- [14] F. L. Roux. Migration des points errants d'homéomorphisme de surface. *C. R. Acad. Sci. Paris*, Série I(t. 330):225–230, 2000.